

- * The chance for children to develop chronic Hepatitis B who were born to mothers with HBeAg positive or Hepatitis B carrier will be reduced by 90% after receiving HBIG and 3 doses of Hepatitis B vaccine. However, there are still 10% chance of becoming a carrier.
- * If the mother is positive for HBsAg, the baby should be injected with **one dose of HBIG and hepatitis B vaccine** as soon as possible after birth. The sooner the better, **not later than 24 hours**.
- * To enable early detection and to understand the effectiveness of the vaccines, patients are tracked and followed up regularly.
- * Starting 2012.09 children born to

mothers with positive HBeAg are tested for HBsAg and anti-HBs at age 12 months. If the babies are tested negative for both, they may receive one dose of Hepatitis B vaccine. 1 month after, their blood will be tested again. If the result for anti-HBs is still negative, the babies may receive second dose after 1 month, and the third dose after 6 months.

- * Mothers who are HBeAg positive should bring the "Hepatitis B record and follow up form for pregnant women" and other information in the maternal health book let to serve as a reference for the pediatrician when examining the child for HBsAg and anti-HBs.



(Picture taken from the Internet)

**Children who were born
to mothers with positive
HBeAg**

**For inquiries: Chung Shan Hospital
Tel: (02) 2708-1166
Nursery Ext.: 2220**

105.06 製、114.07 修

Modern Medicine
Tradition Care